

The Renormalization Group Approach to the Confining String

Enrique Álvarez[‡]¹ and César Gómez[†]²

[‡]*Theory Division, CERN, 1211 Geneva 23, Switzerland,*

^{†‡}*Instituto de Física Teórica, C-XVI,³ and Departamento de Física Teórica, C-XI,
Universidad Autónoma de Madrid E-28049-Madrid, Spain*

Abstract

The renormalization group approach towards the string representation of non abelian gauge theories translates, in terms of the string sigma model beta function equations, the renormalization group evolution of the gauge coupling constant and Zamolodchikov's c function. Tachyon stability, glueball mass gap, renormalization group evolution of the c function and the area law for the Wilson loop are studied for a critical bosonic string vacuum corresponding to a non abelian gauge theory in four dimensional space-time. We prove that the same intrinsic geometry for the string vacuum is universal in some sense, reproducing the Yang-Mills beta function to arbitrary loop order in perturbation theory.

¹E-mail: enrique.alvarez@uam.es, @cern.ch

²E-mail: cesar.gomez@uam.es

³Unidad de Investigación Asociada al Centro de Física Miguel Catalán (C.S.I.C.)

Contents

1	Introduction	2
2	Gauge Geometry	4
2.1	Two loop approximation and universal metric	7
2.2	Generalities on the Confining Manifold	8
2.3	Gravitational Focusing	10
2.4	Other geodesics	11
2.4.1	Null geodesics	12
2.4.2	Timelike geodesics	12
2.4.3	Spacelike geodesics	12
3	Energy functional	13
3.1	Variations of the energy functional	15
4	The propagation of scalar disturbances	18
5	The Wilson loop and the soft dilaton theorem	19

1 Introduction

Phenomenologically the string approach to non abelian gauge theories is based on interpreting the color flux tubes responsible for quark confinement as strings ending on quark-antiquark pairs. From a more theoretical point of view a string representation of a gauge theory amounts to define vacuum expectation values for Wilson loops as sums over random surfaces with boundary given by the loop. After the discovery of D-branes a new way to associate strings with pure gauge theories has been suggested based on strong-weak coupling duality. In that approach, based on Maldacena's conjecture [16], the strong 't Hooft's coupling regime is in correspondence with weakly coupled type IIB string theory. The weakly coupled Yang Mills is in this case described by strongly coupled string theory and therefore out of present technology. A similar problem arises when the Type IIB description is extended to non supersymmetric theories [21] with the supergravity regime being defined by 'tHooft's coupling much larger than the natural cutoff, implying the presence of undesired degrees of freedom.

In the same vein (of working out a string description of the low energy world volume theory on D-branes) type 0 string theories were also considered [12]. There we have a closed string tachyon which means, from the point of view of anti de Sitter space-time, that we should work in a regime where the curvature is of the order of the string scale i.e out of the gravity approximation . Using the vacuum expectation value for the tachyon as source of the dilaton, an approximate solution with a running of the said dilaton was obtained which is consistent with asymptotic freedom .The curvature in the ultraviolet (UV) region being infinity again indicates that the gravity description is not valid in the UV asymptotically free regime. Both in the type 0 case or in type IIB with supersymmetry broken by boundary conditions on the internal Kaluza Klein manifold the starting point was the geometry as defined, in the near horizon limit, by a stack of D branes.

A different approach, presented in a previous paper [5], see also [2], stems from the

idea of using strings to model out the renormalization group equations of the quantum field theory. This in particular leads to associate to the coupling constants of the quantum field theory, some background fields of a closed string theory and to demand that the sigma model beta functions characterizing the string background precisely coincide with the renormalization group equations of the non abelian gauge theory. In this sense the quantum field theory beta functions appear as the input in the string beta function equations. The output, that is purely stringy, will consist in the specific space-time geometry that will be formally defined on the space-time extended by an extra (renormalization group) dimension. This string geometry is the one we can use to define representations of the vacuum expectation value of the Wilson loop in terms of random surfaces, most likely with zig-zag type of boundary conditions [19]. Notice that the difference between the D-brane approach and the renormalization group approach (RGA) lies mainly in the way the field theory input is introduced. In the D-brane approach this is done in terms of the low energy physics on the D-brane. In the RGA approach we use instead the quantum field theory beta functions to fix the dilaton behavior and employ the string beta function equations to derive the stringy space-time geometry.

The physical meaning, from the quantum field theory point of view, of the stringy space-time geometry is related to the renormalization group evolution of Zamolodchikov's c -function. More precisely the curvature on the renormalization group dimension will induce a focusing phenomena with the interpretation of decreasing c -function [22]. In summary the RGA intends to translate in pure gravitational terms, codified in the string beta function equations, the renormalization group evolution of both the gauge coupling constant and Zamolodchikov's c -function. The gravitational description of renormalization group evolution is our way to understand the meaning of holography[2].

We want to continue in this work the study of a critical bosonic string background with the dilaton field defined by an asymptotically free beta function of the type of pure $\mathcal{N} = 0$ Yang-Mills.

The solution presented in reference [5] is generalized in this paper to the case of two loop Yang Mills beta function. The resulting confining metric possesses the same intrinsic geometry that for the one loop case. Moreover the confining geometry is universal for arbitrary order in perturbation theory. This confining string metric is an exact solution to the critical bosonic string sigma model beta function equations for vanishing vacuum expectation value of the tachyon.

Next we consider the problem of stability of our solution showing that on the reduced functional space defined by demanding conservation of energy, the linearized small fluctuation problem does not admit normalizable solutions with negative eigenvalue. We also study the existence of a mass gap for glueball equations as well as the area law behavior for the vacuum expectation value of the Wilson loop. In addition we present a careful analysis of the geometry of the confining string vacuum with the expected behavior of a geometric c -function defined in terms of the expansion for congruences of null geodesics.

2 Gauge Geometry

Our starting point is the RG evolution for an asymptotically free pure gauge theory. For simplicity we will first consider the case of the one loop beta function equation:

$$\mu \frac{g}{d\mu} = -\frac{\beta_0}{(4\pi)^2} g^3 \quad (1)$$

for β some numerical constant depending on the gauge group and g the Yang Mills coupling constant. (The free abelian theory can also be put in this form, with $\beta_0 = 0$). Next we use the standard relation between the gauge coupling constant and the dilaton field:

$$g = e^{\frac{\Phi}{2}} \quad (2)$$

In what follows we will identify the renormalization group coordinate μ with a space-time dimension. From the previous equations we derive the dependence on this coordinate of

the dilaton field Φ :

$$\Phi(\mu) = -\log \log(\mu) + \log(\beta_0) \quad (3)$$

Next we use, in order to fix the string space-time geometry, the sigma model beta function equations for the bosonic string:

$$R_{\mu\nu} + \nabla_\mu \nabla_\nu \Phi = 0 \quad (4)$$

$$(\nabla\Phi)^2 - \nabla^2\Phi = C \quad (5)$$

With the constant C given by:

$$C = \frac{26 - D}{3} \quad (6)$$

To solve these equations for the asymptotically free dilaton field we will use the following ansatz for the metric:

$$ds^2 = a(\mu)d\vec{x}_d^2 + b(\mu)d\mu^2 + c(\mu)d\vec{y}_D^2 \quad (7)$$

(where $d\vec{x}_n^2$ represents the ordinary Minkowski metric in $\mathbb{R}^n$). Before motivating our ansatz for the metric let us just present the solution. Introducing the variable ρ by the relation:

$$\Phi(\mu) = \log(\rho) \quad (8)$$

the metric solution is given by:

$$a(\mu) = \rho \quad (9)$$

$$b(\mu)d\mu^2 = d\rho^2 \quad (10)$$

$$c(\mu) = 1 \quad (11)$$

The corresponding metric is solution provided:

- The number of x coordinates , $d = 4$
- The string is critical i.e the number of y coordinates $D = 21$. and therefore the constant C is equal zero.

- The metric is a solution with both euclidean and Lorentzian signature.

Notice also that the sigma model beta functions we have used assume already a vanishing vacuum expectation value for the closed string tachyon background field.

The metric we obtain as solution to the string beta function equations with an asymptotically free behavior for the dilaton is of the type first considered by Polyakov [19] with the function $a(\mu)$ playing the role of a running string tension. Moreover we can interpret the variable ρ as the analog of Liouville field in non critical strings. It is important to observe that the string beta function equations are not sensitive to the constant in equation (5) i.e they do not feel the concrete gauge group we are working with.

Next we will briefly describe the different regions of our space-time. The region corresponding to ultraviolet behavior of the Yang Mills theory is $\mu = \infty$. In this region the scalar curvature that is given by:

$$R = -\frac{1}{\rho^2} \tag{12}$$

is strong while in the strongly coupled field theory infrared region we have weakly coupled gravity. At the point $\rho = 0$ we have a naked singularity that could be interpreted as the limit of validity of the string description. Comparing with Polyakov's type of metrics we should notice that the singularity satisfy the zig-zag condition $a(\mu = 1) = 0$. Therefore it would be natural to define a string representation of Wilson loops to impose Dirichlet boundary conditions on $\rho = 0$.

As mentioned in the introduction the RGA to the string description of non abelian gauge theories have as its goal to model in terms of sigma model string beta function equations the different renormalization group equations of the corresponding quantum field theory. In this sense it is of major importance to have an interpretation of the string space-time metric in purely quantum field theory terms. Here is where the role of Zamolodchikov's c theorem becomes important. As described in reference [5] the c -theorem can be understood as a direct consequence of gravitational focusing for congruences of null geodesics in the

renormalization group direction μ . Therefore from the RGA point of view the curvature in the coordinate μ should be intimately related with the evolution of c -function. Here we will not enter into a detailed discussion on the definition of the c -function from a geometrical point of view reducing ourselves to discuss the phenomena of gravitational focusing for the asymptotically free space-time metric.

2.1 Two loop approximation and universal metric

Let us now consider the two loop running coupling constant for pure Yang Mills ([18]):

$$g^2(\mu) = \frac{16\pi^2}{\beta_0 \log \mu^2} \left[1 - \frac{2\beta_1}{\beta_0^2} \frac{\log \log \mu^2}{\log \mu^2} + \frac{4\beta_1^2}{\beta_0^4 \log^2 \mu^2} \right. \\ \left. ((\log \log \mu^2 - 1/2)^2 + \frac{\beta_2\beta_0}{8\beta_1^2} - \frac{5}{4}) \right] \quad (13)$$

where the beta function is parametrized as:

$$\beta(g) \equiv \mu \frac{\partial g}{\partial \mu} = -\frac{\beta_0}{(4\pi)^2} g^3 - \frac{2\beta_1}{(4\pi^2)^4} g^5 - \frac{\beta_2}{2(4\pi)^6} g^7 \quad (14)$$

The quotient β_1/β_0^2 happens to be independent, for pure Yang Mills, of the particular gauge group we are working with. (Up to two loops we should neglect β_2). Starting with the three loop contribution, a nontrivial dependence on the renormalization scheme appears.

The corresponding dilaton is given up to a constant by

$$\Phi(\mu) = -\log \log \mu^2 + \log \left[1 - \frac{2\beta_1}{\beta_0^2} \frac{\log \log \mu^2}{\log \mu^2} + \frac{4\beta_1^2}{\beta_0^4 \log^2 \mu^2} \right. \\ \left. ((\log \log \mu^2 - 1/2)^2 + \frac{\beta_2\beta_0}{8\beta_1^2} - \frac{5}{4}) \right] \quad (15)$$

Our ansatz for the confining string metric is now dictated by the one loop solution described in the previous section, namely

$$ds^2 = a(\mu) d\vec{x}_d^2 + b(\mu) d\mu^2 + d\vec{y}_D^2 \quad (16)$$

It is a simple matter (with the help of some algebraic manipulations) to check that the metric:

$$ds^2 = e^\Phi d\vec{x}_d^2 + (de^\Phi)^2 + d\vec{y}_D^2 \quad (17)$$

with Φ the one given by the quantum field theory renormalization group equation, is the desired solution to the sigma model beta function equations. It is important to stress that here also (as already did happen in the one and two loop cases) the sigma model beta function for the dilaton $\beta_\Phi = 0$ implies that the metric (16) is solution only when we consider a four dimensional space-time and a critical string i.e D in (16) equal 21.

At this point the reader can wonder why - if we are identifying the string coupling constant with the Yang Mills coupling- we are using ,in order to match Yang Mills higher loop effects, the sigma model beta function equations at string tree level. This is a very deep question whose answer could be related to the fact that dilaton tadpole effects of Fischler-Susskind type [13] are absorbed in renormalizations of the metric that preserve the universal form (17).

The result we have presented for the two loop case is not accidental, in fact we can prove the following theorem

Theorem. The universal metric (17) is (for an arbitrary dilaton field depending only on the renormalization group variable) an exact solution to the critical string sigma model beta function equations.

This means that *exactly* the same spacetime metric conveys quite different solutions, summarizing the whole perturbative expansion of the gauge theory. A lot of physics is hidden in the change of variables needed to put the metric in the universal form (17).

This is by itself an interesting property of the tree level sigma model beta function equations that depends crucially on working at criticality i.e for the constant (2) vanishing.

2.2 Generalities on the Confining Manifold

In terms of the universal metric (17) it is convenient for most geometrical studies to employ the variable

$$\rho \equiv e^\Phi \tag{18}$$

such that the metric reads

$$ds^2 = \rho d\vec{x}_4^2 + d\rho^2 \quad (19)$$

and the dilaton

$$\Phi = \log \rho \quad (20)$$

The natural range for ρ is $(0, \infty)$ due to the singularity at $\rho = 0$. It is amusing to note that we can extend the metric to the range $(-\infty, 0)$. This is a non-analytic extension, in the sense that the complete metric would read:

$$ds^2 = |\rho| d\vec{x}_4^2 + d\rho^2 \quad (21)$$

and the dilaton

$$\Phi = \log |\rho| \quad (22)$$

This non analytic extension of the metric could be related to the existence of a deconfining phase transition.

We shall use roman capitals to denote generic five-dimensional indices, $A, B, C, \dots = 0, 1, 2, 3, 4$, where $x^4 \equiv z$; and greek letters for the ordinary four-dimensional variables, i.e. $\mu, \nu, \dots = 0, 1, 2, 3$, whereas $i, j, k, \dots = 1, 2, 3$.

The isometry group of the confining manifold is just the four-dimensional Poincaré group, $IO(1, 3)$. The four Killings associated to four-dimensional translations are:

$$k_\mu \equiv \partial_\mu \quad (23)$$

Note in particular that the *contravariant* components of the Killing associated to time translations are

$$k^A \equiv (1, 0, 0, 0, 0) \quad (24)$$

2.3 Gravitational Focusing

The first ingredient we need are the null geodesics for our metric. They are given in affine parametrization by:

$$x^\mu = \frac{2c^\mu\gamma}{c^0} \left(\frac{3c^0}{2\gamma} (\tau - \tau_0) + \rho_0^{3/2} \right)^{1/3} + x_0^\mu - \frac{2c^\mu\gamma\rho_0^{1/2}}{c^0} \quad (25)$$

and

$$\rho = \left(\frac{3c^0}{2\gamma} (\tau - \tau_0) + \rho_0^{3/2} \right)^{2/3} \quad (26)$$

with $\eta_{\mu\nu}c^\mu c^\nu = -\frac{c^0}{\gamma}$.

The covariant null tangent vector is

$$k_\mu = c_\mu \quad (27)$$

(where $c_\mu \equiv \eta_{\mu\nu}c^\nu$) and

$$k_4 = \frac{c^0}{\gamma} \rho^{-1/2} = \frac{c^0}{\gamma} \left(\frac{3c^0}{2\gamma} (\tau - \tau_0) + \rho_0^{3/2} \right)^{-1/3} \quad (28)$$

The affine parameter is normalized in such a way that

$$k^A \nabla_A \tau = 1 \quad (29)$$

The covariant derivative is given by:

$$\begin{aligned} \nabla_\mu k_\nu &= \frac{1}{4} k_4 \eta_{\mu\nu} \\ \nabla_4 k_\mu &= \nabla_\mu k_4 = -\frac{1}{2\rho} k_\mu \\ \nabla_4 k_4 &= -\frac{c^0}{2\gamma\rho^{3/2}} \end{aligned} \quad (30)$$

The fact that the antisymmetric part vanishes is equivalent to the fact that the vorticity of the corresponding congruence $\omega = 0$. This means that the congruence is hypersurface orthogonal.

The expansion is given by

$$\theta = \frac{3c^0}{2\gamma\rho^{3/2}} \quad (31)$$

and the shear by

$$\sigma^2 = -\frac{15c^0}{8\gamma^2\rho^3} \quad (32)$$

We can see directly a c-theorem of sorts operating in the confining metric: the expansion θ behaves in physical variables as

$$\theta \sim (\log \mu)^{3/2} \quad (33)$$

being ∞ at the singularity as it should be. This means that the corresponding c-function as defined in [2] will also have a similar physical behavior. This definition, however was tailored for holographic theories, and it is conceivable that some modifications are needed in our case. It is curious ,however, to notice how easy is to prove this sort of properties in the string representation in contradistinction to the painful attempts to prove it directly by quantum field theory methods [14]. Clearly further research is needed on what quantum field theoretical correlators is the c-function measuring.

2.4 Other geodesics

In order to obtain analytic expressions, it is useful to parametrize the curves in terms of the time coordinate, t . The Minkowskian space-like coordinates have always a form identical to the one corresponding to flat space geodesics, namely:

$$x^i = x_0^i + \beta^i t \quad (34)$$

(where $\beta^i \equiv \frac{c^i}{c^0}$). Let us now examine several subcases in turn.

2.4.1 Null geodesics

Null geodesics exist only when $\vec{\beta}^2 \equiv \beta^2 \leq 1$ (that is, when the Minkowskian projection is timelike). Its explicit form in our present parametrization is:

$$\rho = \frac{1 - \beta^2}{4}(t - t_0)^2 \quad (35)$$

when $\vec{\beta}^2 < 1$, and

$$\rho = \rho_0 \quad (36)$$

when $\vec{\beta}^2 = 1$.

2.4.2 Timelike geodesics

Timelike geodesics also need $\vec{\beta}^2 < 1$. The explicit expression is

$$\rho = \frac{(c^0)^2}{2}(1 - \beta^2)\left(1 + \sin \frac{t - t_0}{c^0 \sqrt{1 - \beta^2}}\right) \quad (37)$$

It is remarkable that this implies the existence of *closed timelike geodesics*; for example, when $\beta^i = 0$:

$$\begin{aligned} x^i &= x_0^i \\ \rho &= \frac{(c^0)^2}{2\gamma^2}\left(1 + \sin \frac{\gamma(t - t_0)}{c^0}\right) \end{aligned} \quad (38)$$

The physical meaning of five-dimensional timelike geodesics is related to unsuspected properties of the renormalization group flow, worth examining in some detail. We stress once more that the ordinary four dimensional projection never suffers from closed timelike curves; they close in the coordinate associated to the renormalization group evolution parameter only.

2.4.3 Spacelike geodesics

Spacelike geodesics appear for any value of β . When $\beta \neq 0$,

$$\rho = -(c^0)^2(1 - \beta^2) \cosh^2 \frac{t - t_0}{c^0 \sqrt{1 - \beta^2}} \quad (39)$$

and when $\beta = 1$,

$$\rho = \rho_0 \pm t \tag{40}$$

3 Energy functional

Since we are considering a background for the critical bosonic string an issue we should address is the fate of tachyons on this particular space-time. Here we will follow the approach of [7] extended from the case of Anti-de-Sitter (AdS) to our confining metric. Let us first recall the AdS argument. We consider the case of a free scalar field of mass m^2 . Next conservation of the energy momentum tensor is enforced. This implies certain boundary conditions on the the fields at the boundary. Finally we solve the field equations on the functional space satisfying these boundary conditions. For stationary wave solutions this amounts to a bound on allowed values for the mass m^2 . Certain negative values of m^2 are allowed by this bound corresponding to tachyons with Compton wavelength of the order of AdS curvature radius. However fluctuations corresponding to these tachyons have positive total energy implying the stability of AdS space-time. In the AdS case the Cauchy problem on the reduced functional space, determined by the condition of energy conservation, has a unique solution. In order to extend the previous analysis to our case the first thing we will do is to derive the boundary conditions on the wave functions required by conservation of energy.

Given an arbitrary scalar field defined on the confining manifold with energy-momentum tensor T_{AB} , there is a *Killing energy* current, namely

$$j^A \equiv T^{AB} k_B \tag{41}$$

which is covariantly conserved,² i.e.

$$\nabla_A j^A = 0 \quad (43)$$

Applying Stokes' theorem to the integral over the five-dimensional region $\mathcal{R}$ delimited by the hypersurfaces $t = t_0$ and $t = t_1$, of the five-form proportional to the first member of (43), i.e. of $d * j$, where j is the one-form dual to the current, $j \equiv j_A dx^A$, we get

$$0 = \int_{\mathcal{M}} d * j = \int_{\partial \mathcal{M}} * j = E(t_2) - E(t_1) + \Delta E \quad (44)$$

where

$$E(t) \equiv \int d^3 x dz j^0(t, x^i, z) \quad (45)$$

and the flux over the boundary at infinity

$$\Delta E = \lim_{(x^i, z) \rightarrow \infty} \int [(\epsilon_{ijk} dt \wedge dx^j \wedge dx^k \wedge dz) j^i + d^4 x j^4] \quad (46)$$

Only when the physical boundary conditions are such that

$$\Delta E = 0 \quad (47)$$

there is actual energy conservation,

$$\frac{dE}{dt} = 0 \quad (48)$$

For a scalar field of mass m it is readily found that

$$T_{AB} = \frac{1}{2} e^\Phi [\nabla_A \phi \nabla_B \phi - \frac{1}{2} g_{AB} (g^{CD} \nabla_C \phi \nabla_D \phi - m^2 \phi^2)] \quad (49)$$

so that the five dimensional energy is:

$$E[\phi] \equiv \frac{1}{4} \int d^3 x dz [\dot{\phi}^2 + (\nabla \phi)^2 + z(m^2 \phi^2 + (\phi')^2)] \quad (50)$$

In the *free* case the energy is zero on shell, provided the fields vanish at infinity fast enough, and this is true no matter what the sign of m^2 is, i.e., even for the putative tachyon.

² Please note that we need at least two scalar fields to build a conserved *number* current, namely

$$N^A \equiv e^\Phi (\phi_2 \partial^A \phi_1 - \phi_1 \partial^A \phi_2) \quad (42)$$

3.1 Variations of the energy functional

Because we are mostly interested in the stability of our solution we will reduce ourselves to study the spectral problem for the operator defined by the second variation of the energy around $T = 0$ corresponding to vanishing tachyonic background. We will use static fluctuations. Notice that the first static variation of the energy is zero since we are working with a solution of the equations of motion.

The second variation can actually be written in our case as:

$$\frac{\delta^2 E[\phi]}{\delta \phi^2} = 2 \int d^3 x dz [-\Delta_3 + L] \quad (51)$$

where L is the operator:

$$L = -z \frac{d^2}{dz^2} - \frac{d}{dz} + zm^2 \quad (52)$$

The issue as to the stability of the solution is then equivalent to the issue of whether the operator L has any negative eigenvalues (on the functional space in which all conservation laws can be actually implemented). We are then led to study the eigenvalue problem

$$LT = \lambda T \quad (53)$$

This is a second order ordinary differential equation, with a regular singular point at the origin and an irregular one at infinity. By studying the indicial equation it is immediately obvious that one of the solutions has a logarithmic singularity at the origin.

In order to be able to study simultaneously the two cases of positive and negative mass square we shall for the latter use $m^2 \equiv -\mu^2$. By performing the change of both dependent and independent variables:

$$T(z) = e^{-z\xi} y(z/\rho) \quad (54)$$

(where $\xi^2 = m^2$ and $2\xi\rho = 1$), the equation is reduced to a confluent hypergeometric one:

$$\frac{d^2 y}{dx^2} + \left(\frac{1}{x} - 1\right) \frac{dy}{dx} + \left(\frac{\lambda}{2m} - \frac{1}{2}\right) \frac{1}{x} y = 0 \quad (55)$$

Let us define $a \equiv \frac{m-\lambda}{2m}$. There is an exceptional case, when $a \in -\mathbb{N}$ (that is, $\lambda = (2p+1)m$, with $p \in \mathbb{Z}^+$), where the solution is given in terms of Laguerre polynomials,

$$y = L_n^{(0)}(x) \quad (56)$$

Otherwise the two independent solutions are given in terms of the functions ([10]) $\phi(a, 1, x)$ and $\psi(a, 1, x)$,

The function $\phi(a, 1, x)$ is regular at the origin

$$\phi(a, 1, x) \equiv {}_1F_1(a, 1, x) \sim_{x \sim 0} 1 + ax + \dots \quad (57)$$

and at infinity behaves as

$$\phi(a, 1, x) \sim_{x \sim \infty} \frac{1}{\Gamma(a)} e^x x^{a-1} \quad (58)$$

On the other hand the function $\psi(a, 1, x)$ has got a logarithmic divergence at the origin

$$\psi(a, 1, x) \sim_{x \sim 0} -\frac{1}{\Gamma(a)} \log[x] + \dots \quad (59)$$

whereas at infinity behaves as

$$\phi(a, 1, x) \sim_{x \sim \infty} x^{-a} \quad (60)$$

When $m^2 > 0$ the only solution regular at the origin is fine. We can arrange our variables in such a way that asymptotically

$$T(z) \sim_{x \sim \infty} e^{-mz} (2mz)^{a-1} \quad (61)$$

This means that the solutions and their derivatives are square integrable on dz .

When $m^2 < 0$, however, the asymptotic behaviour of the putative solution is

$$T \sim e^{\pm i\mu z} (\pm 2i\mu z)^{a-1} \quad (62)$$

(where $a = \frac{1}{2} + \frac{i\lambda}{2\mu}$), too slow to be convergent, for any value of λ .

Thus we can conclude that there is not any negative eigenvalue -even in the tachyonic case $m^2 < 0$ - with normalizable wave function. In other words on the functional space

defined by the boundary conditions required by energy conservation the operator L has not any negative eigenvalue, which we interpret as a strong indication of the stability of our confining metric. Comparing with the bound in [7] we observe that for normalizable static wave functions the mass m^2 is forced to be ≥ 0 . The extension of the previous analysis to stationary wave functions does not change the result.

It is illustrative to present the previous analysis from the point of view of an equivalent Schrodinger problem. In order to do that let us perform the change of variables in equation (53)

$$T(z) = t(z)z^{-\frac{1}{2}} \quad (63)$$

and the equation reduces to

$$\frac{d^2 t}{dz^2} + \left[\frac{\lambda}{z} + \frac{1}{4z^2} - m^2 \right] t = 0, \quad (64)$$

an equation first studied by Whittaker, which is an Schrödinger equation for the reduced potential

$$U(z) = -\frac{1 + 4\lambda z}{4z^2} \quad (65)$$

and reduced energy

$$\epsilon = -m^2 \quad (66)$$

Please note that now we are viewing the problem head down: we are assuming we already know λ , and we look for the energy as an eigenvalue. The potential starts at $U_- = -\infty$ at $z = 0$ can become positive if λ is negative enough, and reaches $U_+ = 0$ at $z = \infty$, but *not faster* than $1/z$ (which is a necessary condition for most inverse scattering theorems to hold)

Again, for $m^2 > 0$, we need the potential to have a negative eigenvalue, which is no problem (there is indeed a continuum spectrum $\epsilon \in (0, -\infty)$).

For $m^2 < 0$, we would need a positive energy, which is not possible with normalizable states, owing to the slow asymptotic behaviour of the potential at infinity.

The massless case is marginal.

4 The propagation of scalar disturbances

From a practical point of view the main utility of a confining string representation of a non-abelian gauge theory ought to be the computation of the glueball spectrum (which is in principle measurable, although the widths are quite big, and the mixing with $q\bar{q}$ states is important).

The main tool in this computation is a calculation of two Wilson loops correlator,

$$\langle W(C)W(C') \rangle \quad (67)$$

which conveys the desired information in the location of its poles. While we certainly plan to perform this computation in the near future, we would like here to examine a very preliminary aspect of this problem, namely the semiclassical propagation of a scalar disturbance in the confining metric (which could be a crude model of the 0^{++} glueball).

We shall consider fields in the four-dimensional momentum representation, i.e.

$$\phi(x, z) \equiv \int d^4k e^{ik \cdot x} \phi(k, z) \quad (68)$$

The wave equation for massless (in the five dimensional sense) scalar disturbances is then

$$z \frac{d^2 \phi}{dz^2} + \frac{d\phi}{dz} = k^2 \phi \quad (69)$$

It is plain to check that there are no normalizable solutions when $k^2 = 0$ (which corresponds to the massless case from the four-dimensional point of view). Actually, the general solution is then

$$\phi(k^2 = 0, z) = c_1 \log z + c_2 \quad (70)$$

This is enough to establish the existence of a *mass gap*, in the sense that massless disturbances from the five-dimensional sense necessarily correspond to massive ones from the four-dimensional one.

When $k^2 \neq 0$ the solution is easily expressed in terms of Bessel functions;

$$\phi(k, z) = B_0(2k\sqrt{z}) \quad (71)$$

where $B_0(x)$ represents a Bessel function; actually a linear combination of the functions $J_0(x)$ and $Y_0(x)$ when $k^2 < 0$ (in this case, the argument in the function should be $2\sqrt{-k^2}\sqrt{z}$), and a linear combination of the modified functions $I_0(x)$ and $K_0(x)$ when $k^2 > 0$.

In the former case there are no normalizable solutions (in accordance with the results in the last paragraph). Physically, this means that there are no *four-dimensional tachyons* propagating. In the latter, normalizability again uniquely selects the K_0 solution.

Normalization now means, for example, that

$$\int_0^\infty \phi(z)^2 dz = 1 \quad (72)$$

But this integral is easily shown to be equivalent to

$$\int_0^\infty K_0^2 z dz = 2k^2 = \frac{1}{2} \quad (73)$$

enforcing a discrete *four dimensional* mass spectrum, namely $m_4^2 = \frac{1}{4}$. Energy is of course measured in units of Λ , so that this gives

$$m = \frac{\Lambda}{2}. \quad (74)$$

This value seems a bit low,³ but of course this is only but a crude estimate and we do not want to push it too far.

5 The Wilson loop and the soft dilaton theorem

An important test of any confining string model is to produce an area law behavior for the vacuum expectation value of the static Wilson loop, which we shall take in the static

³ Experimentally the situation is unclear; although in [18] several candidates are reported, notably the $f_0(1500)$. In [11] evidence is reported for a Pomeron trajectory $\alpha_P(t) = \alpha_P(0) + \alpha'_P t$, with $\alpha_P(0) = 1.08 GeV$ and $\alpha'_P = 0.25 GeV^{-2}$. The standard lattice value (which seems to be a decreasing function of the N factor in the gauge group $SU(N)$), is about $m \sim 3.64\sqrt{\sigma} \sim 2.18 GeV$ [20]. In order to compare with the string value, an extrapolation towards $N = \infty$ ought to be made.

gauge

$$\begin{aligned}x^0 &= \tau \\ x^1 &= \sigma\end{aligned}\tag{75}$$

and the world sheet of the confining string will be characterized by the time translation of the profile

$$\rho = \rho(\sigma)\tag{76}$$

with $\rho(\pm l/2) = 0$. Once we have a string model of a pure gauge theory we should represent the vacuum expectation value for the Wilson loop as a sum over random surfaces with fixed boundary as defined by the loop. In the semiclassical approximation we can use the corresponding Nambu-Goto form of the string action and to look for the minimal area surface bounded by the loop [17].

For the confining string metric the Nambu Goto action is given by:

$$S = \frac{1}{2\pi\alpha'} \int \sqrt{\rho((\partial_x \rho)^2 + \rho)}\tag{77}$$

where by simplicity we use a static loop with profile given in terms of the function $\rho(x)$. Denoting ρ_0 the minimum value of $\rho(x)$ at the symmetric point $x = 0$ it is easy to see that this value depends on L , the spatial length of the loop, as $\rho_0 \sim L^2$.

If we use in the Nambu Goto action a constant value for the string tension α' the potential energy in this semiclassical approximation will go as:

$$V(L) \sim L^3\tag{78}$$

which means overconfining. It is important to stress that this result is generic for our metric independently if we locate the loop at $\rho = 0$ or at $\rho = \text{infy}$. The only difference is that when located in the zig-zag hyperplane $\rho = 0$ the minimal surface will live in the non analytic extension of the space corresponding to negative values of ρ (i.e to values of μ between zero and one).

An important theorem for the bosonic string is the so called soft dilaton theorem that establish the following interesting relation between the string coupling constant and the string tension:

$$\lim_{k \rightarrow 0} A(k; p_1, p_2, \dots p_n) = cg(\sqrt{\alpha'})^{\frac{(d-2)}{2}} (\sqrt{\alpha'} \frac{\partial}{\partial \sqrt{\alpha'}} - \frac{1}{2}(d-2)g \frac{\partial}{\partial g}) A(p_1, p_2, \dots p_n) \quad (79)$$

Where $A(k; p_1, p_2, \dots p_n)$ is the amplitude for a soft dilaton insertion of momentum k and n external gluon vertex operators with momentum $(p_1, p_2, \dots p_n)$. If now we identify the open string coupling constant g with the Yang Mills coupling, something that we have done along our construction then the soft dilaton theorem [1] will implies that if g is running with the renormalization group parameter μ then $\sqrt{\alpha'}^{\frac{(d-2)}{2}}$ should change with μ in precisely the opposite way. Notice also that in the soft dilaton theorem expression we must consider the momentum $(p_1, p_2, \dots p_n)$ living in the four dimensional physical space-time. This is necessary in order to be consistent with our identification of the string coupling constant with the gauge coupling thus we will take $d = 4$.

The way we will use now the soft dilaton theorem is by simply changing the string tension in the Nambu-Goto action by the effective running string tension dictated by the soft dilaton theorem relation between the scaling of g and α' . In these conditions we get:

$$S = \frac{1}{2\pi} \int \sqrt{\rho^{-1}((\partial_x \rho)^2 + \rho)} \quad (80)$$

It is now plain to see that the potential energy in this case goes like

$$V(L) \sim L \quad (81)$$

i.e as the area law. The reader can wonder what will happen if instead of taking $d = 4$ in the soft dilaton theorem we leave free this value. The potential we will get is given in terms of d as

$$V(L) \sim L^{5-d} \quad (82)$$

producing the area law only in the four dimensional case.

The reader should notice that the Wilson loop is ,properly speaking, the first string computation we are performing. Before we were considering semiclassical gravity effects associated with our confining string metric. Very likely the way the soft dilaton theorem enters into the computation of the Wilson loop is related to loop equations understood as renormalization group equations for the vacuum expectation value of the Wilson loop. It is also important to stress that the soft dilaton theorem ensures the area law behavior for the universal confining metric (17) whatever the loop order of the Yang Mills beta function we choose to fix the dilaton running.

There is still a problem we have not touched along our paper, that can be relevant for our discussion of the Wilson loop computation, namely the dynamical role of the 21 spectator extra coordinates we need to add in order to find solution to the sigma model beta function equations. We have not a definitive answer to this issue although we feel very meaningful the fact that our solution make it compelling to work in a critical string framework. (incidentally notice that this would be also the case even in the type 0 strings setup with world sheet supersymmetry). A potential way to interpret these extra spectators is to work in non critical dimension with an appropriate vacuum expectation value for the tachyon. In this case, and in order to solve the tachyon beta function equation we need to invoke some mechanism of the type described in reference ([12]) for the generation of a tachyon potential $V(T)$ such that $mT_0 + V'(T_0) = 0$ where T_0 is the required tachyon vacuum expectation value. In spite of the fact that this can be an interesting mechanism to avoid the presence of extra spectators we insist, until reaching a deeper understanding on the meaning of criticality, on the beauty of having an exact string solution with a running dilaton consistent with higher order effects in quantum field perturbation theory.

Acknowledgments

We are grateful for stimulating discussions with Luis Alvarez-Gaumé, Peter Hasenfratz and Peter Minkowski. We are indebted to Gabriele Veneziano for pointing out an error in an earlier version of the manuscript. This work has been partially supported by the European Union TMR program FMRX-CT96-0012 *Integrability, Non-perturbative Effects, and Symmetry in Quantum Field Theory* and by the Spanish grant AEN96-1655. The work of E.A. has also been supported by the European Union TMR program ERBFMRX-CT96-0090 *Beyond the Standard model* and the Spanish grant AEN96-1664.

References

- [1] M. Ademollo, A. D’Adda, R. D’Auria, F. Gliozzi, E. Napolitano, S. Sciuto, P. Di Vecchia *Soft Dilatons and scale renormalization in dual theories*. Nucl.Phys.B94:221,1975
- [2] E.T.Akhmedov, *A Remark on the AdS/CFT correspondence and the Renormalization Group Flow*, hep-th/9806217, Phys.Lett. B442 (1998) 152;
E. Alvarez and C. Gómez, *Geometric Holography, the Renormalization Group and the c-Theorem*, hep-th/9807226; Nucl. Phys. B541 (1999),441;
V. Balasubramanian and P. Kraus, *Spacetime and the Holographic Renormalization Group* hep-th/9903190 ;
D. Z. Freedman, S. S. Gubser, K. Pilch, N. P. Warner, *Renormalization Group Flows from Holography– Supersymmetry and a c-Theorem*, hep-th/9904017;
Vatche Sahakian, *Holography, a covariant c-function, and the geometry of the renormalization group*, hep-th/9910099.
- [3] O. Alvarez, *The static potential in string models*, Phys. Rev, D24,440 (1981)

- [4] E. Alvarez and C. Gómez, *Non-critical Confining Strings and the Renormalization Group*, hep-th/9902012; Nucl. Phys. B550 (1999), 169.
- [5] E. Alvarez and C. Gómez, *The Confining String from the Soft Dilaton Theorem* hep-th/9907158, Nucl. Phys, to be published.
- [6] E. Alvarez, C. Gomez and T. Ortin, *String Representation of Wilson Loops*, . hep-th/9806075, Nucl.Phys. B545 (1999) 217-232.
- [7] P. Breitenlohner and D. Freedman. *Stability in gauge extended supergravity*, Ann Phys. 144 (1982), 249.
- [8] C.G.Callan, E.J. Martinec, M.J. Perry , D. Friedan , *Strings in background fields*, Nucl.Phys.B262:593,1985
- [9] G. Curci and G. Paffuti, *Consistency between the string background field equation of motion and the vanishing of the conformal anomaly*, Nucl.Phys.B286:399,1987
- [10] A. Erdelyi et al, *Higher trascendental functions*, vol 1 (MacGraw Hill,1954)
- [11] J. Forshaw and D.A. Ross, *QCD and the Pomeron*, (Cambridge).
- [12] I. Klebanov and A. Tseytlin, *D-Branes and Dual Gauge Theories in Type 0 Strings*, Nucl.Phys.B546:155-181,1999 hep-th/9811035
- [13] W. Fischler and L.Susskind *Dilaton Tadpoles, String Condensates and Scale Invariance*. Phys.Lett.B173:262,1986
- [14] S. Forte and Latorre, *A proof of the irreversibility of renormalization group flows in four dimensions*, Nucl.Phys. B535 (1998) 709-728;
H. Osborn and G. Shore, *Correlation Functions of the Energy Momentum Tensor on Spaces of Constant Curvature*, hep-th/9909043.

- [15] M. Lüscher, K. Symanzik and P. Weisz, *Anomalies in the free loop wave equation in the WKB approximation*, Nucl. Phys. B173(1980) 365.
- [16] J. Maldacena, *The large N limit of superconformal field theories and supergravity*, Adv.Theor.Math.Phys.2:231-252,1998, [hep-th/9711200](#).
- [17] J. Maldacena, *Wilson loops in large N field theories*, [hep-th/9803002](#), Phys.Rev.Lett. 80 (1998) 4859-4862
Soo-Jong Rey, Jungtae Yee, *Macroscopic Strings as Heavy Quarks of Large N Gauge Theory and Anti-de Sitter Supergravity*, [hep-th/9803001](#)
- [18] Particle Data Group, *Review of Particle Physics*, Eur. Phys. J. C 3,1 (1998).
- [19] A.M. Polyakov, *The Wall of the Cave*, [hep-th/9808057](#)
- [20] Michael J. Teper, *Gleball masses and other physical properties of $SU(N)$ gauge theories in $D=3+1$: a review of lattice results for theorists*, [hep-th/9812187](#)
- [21] E. Witten, *Anti-de-Sitter space, thermal phase transition and confinement in gauge theories*, [hep-th/9803131](#)
- [22] A. B. Zamolodchikov, *Irreversibility of the flux of the renormalization group in a 2d field theory*, JETP Lett, 43 (1986) 730.